

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
18 July 2002 (18.07.2002)

PCT

(10) International Publication Number  
**WO 02/055502 A1**

(51) International Patent Classification<sup>7</sup>: C07D 213/85,  
213/82, 213/84, 213/61, 213/48, 213/26, 213/38, 213/40,  
A61K 31/4418, A61P 25/00, 29/00

(21) International Application Number: PCT/JP01/11241

(22) International Filing Date:  
21 December 2001 (21.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
PR2372 2 January 2001 (02.01.2001) AU  
PR3007 9 February 2001 (09.02.2001) AU  
PR6697 30 July 2001 (30.07.2001) AU

(71) Applicant (for all designated States except US): FUJISAWA PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ISHIDA, Junya [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). YAMAMOTO, Hirofumi [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). KONISHI, Nobukiyo [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). MORITA, Masataka [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). NAKAMURA, Katsuya [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). MIYATA, Susumu [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP).

OCHI, Takehiro [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). MORITA, Yoshiaki [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). YOSHIMI, Eiji [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP). KURODA, Kanae [JP/JP]; Fujisawa Pharmaceutical Co., Ltd., 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP).

(74) Agent: TABUSHI, Eiji; Fujisawa Pharmaceutical Co., Ltd., Osaka Factory, 1-6, Kashima 2-chome, Yodogawa-ku, Osaka-shi, Osaka 532-8514 (JP).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRIDINE DERIVATIVES USEFUL AS CYCLOOXYGENASE INHIBITOR



(57) Abstract: A compound of the formula (I), wherein R<sup>1</sup> is hydrogen, halogen, carbymoyl, cyano, formyl, or lower alkyl optionally substituted with halogen, amino or a protected amino; R<sup>2</sup> is hydrogen, halogen, cyano or lower alkoxy; R<sup>3</sup> is phenyl or pyridyl, each of which is substituted with lower alkoxy; and R<sup>4</sup> is lower alkoxy; provided that either R<sup>1</sup> or R<sup>2</sup> is hydrogen, then the other is other than hydrogen, or its salts, which are useful as a medicament.

## DESCRIPTION

## PYRIDINE DERIVATIVES USEFUL AS CYCLOOXYGENASE INHIBITORS

5     Technical Field

This invention relates to novel pyridine compounds having pharmacological activity, to a process for their production and to a pharmaceutical composition containing the same.

10    Background Art

The presence of two cyclooxygenase isoenzymes, cyclooxygenase-I (COX-I) and cyclooxygenase-II (COX-II) is known (Proc. Nat. Acad. Sci. USA 88, 2692-2696 (1991)).

Traditional non steroidal anti-inflammatory compounds (NSAIDs) have inhibiting activities of both COX-I and COX-II (J. Biol. Chem., 268, 6610-6614 (1993), etc). The therapeutic use thereof involves undesired effects on the gastrointestinal tract, such as bleeding, erosions, gastric and intestinal ulcers, etc.

It was reported that selective inhibition of COX-II shows anti-inflammatory and analgesic activities comparable with conventional NSAIDs but with a lower incidence of some gastrointestinal undesired effects (Proc. Nat. Acad. Sci. USA, 91, 3228-3232 (1994)). Accordingly, various selective COX-II inhibitors have been prepared. However, it was reported that those "selective COX-II inhibitor" show some side-effects on kidney and/or insufficient efficacy on acute pains.

Further, some compounds such as SC-560, mofezolac, etc, which have certain selective inhibiting activity against COX-I. WO98/57910 shows some compounds having such activity. However, their selectivity of inhibiting COX -I does not seem to be enough to use them as a clinically acceptable and satisfactory analgesic agent due to their gastrointestinal disorders.

And further, some pyridine derivatives having cyclooxygenase-II inhibiting activity have already been known by WO96/24584 and WO98/03484.

## Disclosure of Invention

This invention relates to pyridine compounds, which have pharmaceutical activity such as cyclooxygenase (hereinafter described as COX) inhibiting activity, to a process for their production, to a pharmaceutical composition containing the same and to a use thereof.

Accordingly, one object of this invention is to provide the pyridine compounds, which have a COX inhibiting activity.


Another object of this invention is to provide a process for production of the pyridine compounds.

A further object of this invention is to provide a pharmaceutical composition containing, as active ingredients, the pyridine compounds.

Still further object of this invention is to provide a use of the pyridine compounds for manufacturing a medicament for treating or preventing various diseases.

The new pyridine compounds of this invention can be represented by the following general formula (I):

20



wherein R<sup>1</sup> is hydrogen, halogen, carbamoyl, cyano, formyl, or lower alkyl optionally substituted with halogen, amino or a protected amino;

R<sup>2</sup> is hydrogen, halogen, cyano or lower alkoxy;

R<sup>3</sup> is phenyl or pyridyl, each of which is substituted with lower alkoxy; and


R<sup>4</sup> is lower alkoxy;

provided that either R<sup>1</sup> or R<sup>2</sup> is hydrogen, then the other

is other than hydrogen,  
or its salts.

The compounds (I) or its salts are able to be produced in  
a similar manner to the general processes and Examples shown below.

Process 1




or a salt thereof

or a salt thereof

, in which  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  are each as defined above, and  $R^5$  is  
a leaving group, such as halogen, trifluoromethanesulfonyloxy,  
etc.



(to be continued to the next page)

Process 2




(III)

or a salt thereof




(IV)

or a salt thereof




(Ib)

or a salt thereof




(Ia)

or a salt thereof



(Ic)

or a salt thereof



(Id)

or a salt thereof

, in which  $\text{R}^3$  and  $\text{R}^4$  are each as defined above, and  
X is halogen.

The compounds of formula (I) may contain one or more asymmetric centers and thus they can exist as enantiomers or diastereoisomers. This invention includes both mixtures and separate individual isomers.

5 The compounds of the formula (I) may also exist in tautomeric forms and the invention includes both mixtures and separate individual tautomers.

10 The compounds of the formula (I) and its salts can be in a form of a solvate, which is included within the scope of the present invention. The solvate preferably include a hydrate and an ethanolate.

Also included in the scope of invention are radiolabelled derivatives of compounds of formula (I) which are suitable for biological studies.

15 In the above and subsequent description of the present specification, suitable examples of the various definitions to be included within the scope of the invention are explained in detail in the following.

20 The term "lower" is intended to mean a group having 1 to 6 carbon atom(s), unless otherwise provided.

Suitable "lower alkyl" and lower alkyl moiety in the term "lower alkoxy" may be a straight or branched one, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl, or the like, in which preferable one is methyl.

Suitable "halogen" may be fluoro, chloro, bromo or iodo or the like, which preferable one is chloro.

30 Suitable amino-protective group in the term "a protected amino" is acyl (such as, lower alkanoyl, carbamoyl, etc), lower alkyl, etc.

35 Suitable "lower alkyl optionally substituted with halogen, amino or a protected amino" is lower alkyl; lower alkyl substituted

with halogen; lower alkyl substituted with amino; or lower alkyl substituted with a protected amino.

More preferable "lower alkyl substituted with halogen" is difluoromethyl, trifluoromethyl, or the like, in which the most preferable one is difluoromethyl.

More preferable "lower alkyl substituted with amino" is aminomethyl, aminoethyl, or the like, in which the most preferable one is aminomethyl.

More preferable "lower alkyl substituted with a protected amino" is mono- or di-lower alkylamino(lower)alkyl, such as methylaminomethyl, dimethylaminomethyl, etc; lower alkanoylamino(lower)alkyl, such as acetylaminomethyl; methylcarbamoylaminoethyl.

Suitable "phenyl or pyridyl, each of which is substituted with lower alkoxy" is 4-(lower)alkoxyphenyl or 6-(lower)alkoxy-pyridine-3-yl, in which suitable lower alkyl moiety may be the same as the before-mentioned lower alkyl. The most preferable one is 4-methoxyphenyl or 6-methoxypyridine-3-yl.

Among the compound (I), the following compounds are exemplified as the more preferable ones.

1) The compound, in which

$R^1$  is halogen, carbamoyl, cyano, formyl, or lower alkyl optionally substituted with halogen, amino or a protected amino;

$R^2$  is hydrogen;

$R^3$  is phenyl substituted with lower alkoxy, or pyridyl substituted with lower alkyl; and

$R^4$  is lower alkoxy.

2) The compound, in which

$R^1$  is hydrogen;

$R^2$  is halogen, cyano or lower alkoxy;  
 $R^3$  is phenyl substituted with lower alkoxy; and  
 $R^4$  is lower alkoxy.

- 5           3)    The compound, in which  
                $R^1$  is cyano, or lower alkyl optionally substituted with  
                   halogen, amino or a protected amino;  
                $R^2$  is halogen, cyano or lower alkoxy;  
                $R^3$  is phenyl substituted with lower alkoxy; and  
 10            $R^4$  is lower alkoxy.

Suitable salts of the compounds (I) are pharmaceutically acceptable conventional non-toxic salts and include a metal salt such as an alkali metal salt (e.g., sodium salt, potassium salt, etc.) and an alkaline earth metal salt (e.g., calcium salt, magnesium salt, etc.), an ammonium salt, an organic base salt (e.g., trimethylamine salt, triethylamine salt, pyridine salt, picoline salt, dicyclohexylamine salt, etc.), an organic acid salt (e.g., acetate, maleate, tartrate, methanesulfonate, benzenesulfonate, formate, toluenesulfonate, trifluoroacetate, etc.), an inorganic acid salt (e.g., hydrochloride, hydrobromide, sulfate, phosphate, etc.), a salt with an amino acid (e.g., arginine, aspartic acid, glutamic acid, etc.), or the like.

25           In order to illustrate the usefulness of the object compounds (I), the pharmacological test data of the compounds (I) are shown in the following.

[A]   ANALGESIC ACTIVITY :

30           Effect on adjuvant arthritis in rats :

(i) Test Method :

Arthritis was induced by injection of 0.5 mg of Mycobacterium tuberculosis (Difco Laboratories, Detroit, Mich.) in 50  $\mu$ l of liquid paraffin into the right hind footpad of Lewis rats aged



7 weeks. Analgesic activity of a single dose of agents in arthritic rats was studied. Arthritic rats were randomized and grouped (n=10) for drug treatment based on pain threshold of left hind paws and body weight on day 22. Drugs (Test compounds) were administered and the pain threshold was measured 2hr after drug administration. The intensity of hyperalgesia was assessed by the method of Randall - Selitto. The mechanical pain threshold of the left hind paw (uninjected hind paw) was determined by compressing the ankle joint with a balance pressure apparatus (Ugo Basile Co. Ltd., Varese, Italy). The threshold pressure of rats squeaking or struggling was expressed in grams. The threshold pressure of rats treated with drugs was compared with that of non-treated rats. A dose showing the ratio of 1.5 is considered to be the effective dose.

(ii) Test Results :

Test compound (Example No.)	Dose (mg/kg)	The coefficient of analgesic
Example 4	3.2	> 1.5
Example 13-(5)	3.2	> 1.5

[B] Inhibiting activity against COX-I and COX-II

(Whole Blood Assay):

(i) Test Method :

Whole blood assay for COX-I

Fresh blood was collected by syringe without anticoagulants from volunteers with consent. The subjects had no apparent inflammatory conditions and had not taken any medication for at least 7 days prior to blood collection. 500  $\mu$ l aliquots of human whole blood were immediately incubated with 2  $\mu$ l of either dimethyl sulfoxide vehicle or a test compound at final concentrations for 1hr at 37°C to allow the blood to clot. Appropriate treatments (no incubation) were used as blanks. At the end of the incubation, 5  $\mu$ l of 250mM Indomethacin was added to stop the reaction. The

blood was centrifuged at 6000 x g for 5min at 4°C to obtain serum. A 100  $\mu$ l aliquot of serum was mixed with 400  $\mu$ l methanol for protein precipitation. The supernatant was obtained by centrifuging at 6000 x g for 5min at 4°C and was assayed for TXB<sub>2</sub> using an enzyme immunoassay kit according to the manufacturer's procedure. For a test compound, the results were expressed as percent inhibition of thromboxane B<sub>2</sub> (TXB<sub>2</sub>) production relative to control incubations containing dimethyl sulfoxide vehicle. The data were analyzed by that a test compound at the indicated concentrations was changed log value and was applied simple linear regression. IC<sub>50</sub> value was calculated by least squares method.

#### Whole blood assay for COX-II

Fresh blood was collected in heparinized tubes by syringe from volunteers with consent. The subjects had no apparent inflammatory conditions and had not taken any medication for at least 7 days prior to blood collection. 500  $\mu$ l aliquots of human whole blood were incubated with either 2  $\mu$ l dimethyl sulfoxide vehicle or 2  $\mu$ l of a test compound at final concentrations for 15min at 37°C. This was followed by incubation of the blood with 10  $\mu$ l of 5mg/ml lipopolysaccharide for 24hr at 37°C for induction of COX-II. Appropriate PBS treatments (no LPS) were used as blanks. At the end of the incubation, the blood was centrifuged at 6000 x g for 5min at 4°C to obtain plasma. A 100  $\mu$ l aliquot of plasma was mixed with 400  $\mu$ l methanol for protein precipitation. The supernatant was obtained by centrifuging at 6000 x g for 5min at 4°C and was assayed for prostagrandin E<sub>2</sub> (PGE<sub>2</sub>) using a radioimmunoassay kit after conversion of PGE<sub>2</sub> to its methyl oximate derivative according to the manufacturer's procedure. For a test compound, the results were expressed as percent inhibition of PGE<sub>2</sub> production relative to control incubations containing dimethyl sulfoxide vehicle. The data were analyzed by that a test compound at the indicated concentrations was changed log value

and was applied simple linear regression. IC<sub>50</sub> value was calculated by least squares method.

(ii) Test Results :

Test Compound (Example No.)	COX-I IC <sub>50</sub> (μM)	COX-II IC <sub>50</sub> (μM)
Example 1-(3)	< 0.1	> 1
Example 4	< 0.1	> 1
Example 6	< 0.1	> 1
Example 11-(4)	< 0.1	> 1

It appeared, from the above-mentioned Test Results, that the compound (I) or pharmaceutically acceptable salts thereof of the present invention have an inhibiting activity against COX, particularly a selective inhibiting activity against COX-I.

Additionally, it was further confirmed that the compounds (I) of the present invention lack undesired side-effects of non-selective NSAIDs, such as gastrointestinal disorders, bleeding, renal toxicity, cardiovascular affection, etc.

The object compound (I) or pharmaceutically acceptable salts thereof of this invention possesses COX inhibiting activity and possesses strong anti-inflammatory, antipyretic, analgesic, antithrombotic, anti-cancer activities, and so on. The object compound (I) and pharmaceutically acceptable salt thereof, therefore, are useful for treating and/or preventing COX mediated diseases, inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunological diseases, thrombosis, cancer and neurodegenerative diseases in human beings or animals by using administered systemically or topically. More particularly, the object compound (I) and pharmaceutically acceptable salts thereof are useful for treating and/or preventing inflammation and acute or chronic pain in joint and muscle [e.g. rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis,

gouty arthritis, juvenile arthritis, etc.], inflammatory skin condition [e.g. sunburn, burns, eczema, dermatitis, etc.], inflammatory eye condition [e.g. conjunctivitis, etc.], lung disorder in which inflammation is involved [e.g. asthma, bronchitis, pigeon fancier's disease, farmer's lung, etc.], condition of the gastrointestinal tract associated with inflammation [e.g. aphthous ulcer, Chrohn's disease, atopic gastritis, gastritis varialoforme, ulcerative colitis, coeliac disease, regional ileitis, irritable bowel syndrome, etc.], gingivitis, inflammation, pain and tumescence after operation or injury, pyrexia, pain and other conditions associated with inflammation, particularly those in which lipooxygenase and cyclooxygenase products are a factor, systemic lupus erythematosus, scleroderma, polymyositis, tendinitis, bursitis, periarteritis nodosa, rheumatic fever, Sjogren's syndrome, Behcet disease, thyroiditis, type I diabetes, nephrotic syndrome, aplastic anemia, myasthenia gravis, uveitis contact dermatitis, psoriasis, Kawasaki disease, sarcoidosis, Hodgkin's disease, Alzheimers disease, or the like. Additionally, the object compound (I) or a salt thereof is expected to be useful as therapeutical and/or preventive agents for cardiovascular or cerebrovascular diseases, the diseases caused by hyperglycemia and hyperlipemia.

For therapeutic purpose, the compound (I) and a pharmaceutically acceptable salt thereof of the present invention can be used in a form of pharmaceutical preparation containing one of said compounds as an active ingredient, in admixture with a pharmaceutically acceptable carrier such as an organic or inorganic solid or liquid excipient suitable for oral, parenteral or external administration. The pharmaceutical preparations may be capsules, tablets, dragees, granules, inhalant, suppositories, solution, lotion, suspension, emulsion, ointment, gel, or the like. If desired, there may be included in these preparations, auxiliary substances, stabilizing agents, wetting or emulsifying agents, buffers and other commonly used additives.

While the dosage of therapeutically effective amount of the compound (I) will vary depending upon the age and condition of each individual patient, an average single dose of about 0.01 mg, 0.1 mg, 1 mg, 10 mg, 50 mg, 100 mg, 250 mg, 500 mg and 1000 mg of the compound (I) may be effective for treating the above-mentioned diseases. In general, amounts between 0.01 mg/body and about 1,000 mg/body may be administered per day.

And further, it was also confirmed that analgesic agent is acceptable and satisfactory to patients if its selectivity of inhibiting activity against COX-I, i.e., cyclooxygenase-II vs. cyclooxygenase-I  $IC_{50}$  values ratio ( $IC_{50}$  against COX-II/ $IC_{50}$  against COX-I) is higher than 30 in a whole blood assay, due to a lack of undesired side effects, such as, gastrointestinal disorders, bleeding, renal toxicity, cardiovascular affection, etc. Until now, no one could know what kind of selectivity should be achieved for producing a clinically acceptable and satisfactory "selective COX-I inhibitor" and no one could produce such kind of "selective COX-I inhibitor".

Accordingly, one object of this invention is to provide an analgesic agent comprising a selective cyclooxygenase-I inhibitor, a cyclooxygenase-II vs. cyclooxygenase-I  $IC_{50}$  values ratio of which is higher than 30 in a whole blood assay. More preferable selectivity thereof is higher than 50, and most preferable one is higher than 100.

The selectivity of cyclooxygenase-I inhibitors can be determined by analyzing their  $IC_{50}$  values against cyclooxygenase-II and cyclooxygenase-I in a whole blood assay and by calculating  $IC_{50}$  values ratio thereof.

In the present invention, the "whole blood assay" means an assay method by using whole blood, particularly human whole blood. The inhibiting activity of test compounds against COX-I can be confirmed by assaying the inhibition of  $TXB_2$  production in a human whole blood. And the inhibiting activity of test compounds against

COX-II can be confirmed by assaying the inhibition of PGE<sub>2</sub> in a human whole blood.

Details thereof are shown by "[B] Inhibiting activity against COX-I and COX-II" in the present application. And, the selectivity of test compounds against COX-I and COX-II can be confirmed thereby.

In addition to the above IC<sub>50</sub> values ratio, it is preferable that the cyclooxygenase-II IC<sub>50</sub> value of "selective cyclooxygenase-I inhibitor" is higher than 0.2  $\mu$ M in whole blood assay, more preferably higher than 0.5  $\mu$ M, and most preferably higher than 1.0  $\mu$ M, in order to remove the effect of COX-II inhibiting activity.

The present invention also provides a method for selecting a cyclooxygenase-I inhibitor that lacks gastrointestinal disorders, by assessing whether cyclooxygenase-II vs. cyclooxygenase-I IC<sub>50</sub> values ratio is higher than 30, more preferably higher than 50, and most preferably higher than 100, in whole blood assay.

In order to prove the above invention in more details, the following pharmacological data are shown.

[1] Selective inhibiting activity against COX-I in whole blood assay

IC<sub>50</sub> values of various test compounds were obtained according to a similar manner to the test method shown in "[B] Inhibiting activity against COX-I and COX-II" described in the above. And their selectivity against COX-I was assessed by calculating cyclooxygenase-II vs. cyclooxygenase-I IC<sub>50</sub> values ratio. The results are shown in Table 1.

[2] ANALGESIC ACTIVITY :

Effect on adjuvant arthritis in rats :

Test Method (the same as [A]):

Arthritis was induced by injection of 0.5 mg of Mycobacterium tuberculosis (Difco Laboratories, Detroit, Mich.) in 50  $\mu$ l of liquid paraffin into the right hind footpad of Lewis rats aged 7 weeks. Analgesic activity of a single dose of agents in arthritic rats was studied. Arthritic rats were randomized and grouped (n=10) for drug treatment based on pain threshold of left hind paws and body weight on day 22. Drugs (Test compounds) were administered and the pain threshold was measured 2hr after drug administration. The intensity of hyperalgesia was assessed by the method of Randall - Selitto. The mechanical pain threshold of the left hind paw (uninjected hind paw) was determined by compressing the ankle joint with a balance pressure apparatus (Ugo Basile Co. Ltd., Varese, Italy). The threshold pressure of rats squeaking or struggling was expressed in grams. The threshold pressure of rats treated with drugs was compared with that of non-treated rats. The ratio was shown in Table 1. A dose showing the ratio of 1.5 is considered to be the effective dose.

[3] Stretching test:

Male ddY mice were used after a 24 h fast. Drugs were orally administered to groups of 10 mice. Mice were injected intraperitoneally (i.p.) with 0.2ml/10g of 0.6% acetic acid 1 h after the drug administration and then placed singly in a plastic animal cage. Stretching responses, defined as constriction of the abdomen with stretching of the hind limbs, were counted for 10 min from 3 min after the i.p. injection of acetic acid. The results are shown in Table 1.

[4] Gastric ulcerogenic activity in rats:

Male Sprague-Dawley rats were used after a 24 h fast. Drugs were orally administered to groups of 10 rats 5 h before autopsy. The stomachs were macroscopically inspected and scored as follows: 0, no evidence of gastric lesions; 1, spotty submucosal hemorrhage

or appearance of erosion; 3, widespread adherence of blood and large areas of submucosal hemorrhage or one to four small ulcers; 4, more than four small ulcers or one large ulcer; 5, numerous large ulcers. The results are shown in Table 1.

5

Table 1

Test compound	Whole blood assay (IC <sub>50</sub> : $\mu$ M)			Analgesic activity in adjuvant arthritis -3.2mg/Kg-	Inhibiting activity in stretching test	Gastric ulcerogenic activity  (Non-toxic dose)
	COX-I	COX-II	Selectivity			
A	0.011	8.5	770	1.62	68% (32mg/Kg)	> 100 mg/Kg
B	0.015	3.8	253	1.54	62% (32mg/Kg)	> 100 mg/Kg
C	0.017	1.9	112	1.57	48% (10mg/Kg)	> 100 mg/Kg
D	0.012	0.65	54	1.59	58% (10mg/Kg)	> 100 mg/Kg
References	E	0.0024	0.10	42	-	< 100
	F	0.011	0.15	14	-	< 100
	G	0.054	0.21	3.9	-	3.2
	H	0.42	0.63	1.5	-	10
	I	0.18	0.19	1.1	-	1
	J	0.15	0.028	0.19	-	3.2

("C": Compound produced by Example 4 in the present invention,  
 "F": SC-560, "G": Ketoprofen, "H": Mofezolac,  
 "I": Indomethacin, "J": Diclofenac)

10

From the above experimental data, SC-560 and Ketoprofen, still show insufficient selectivity of inhibiting activity against COX-I and thereby gastrointestinal disorders, though they are announced as "selective COX-I inhibitor" in general. And it was confirmed that the selectivity of inhibiting activity against

15



COX-I, i.e.,  $IC_{50}$  values ratio, should be more than 30, and that cyclooxygenase-II  $IC_{50}$  value should be higher than  $0.2 \mu M$  in whole blood assay.

In other word, the selective cyclooxygenase-I inhibitor, that (1) has a cyclooxygenase-II vs. cyclooxygenase-I  $IC_{50}$  values ratio higher than 30 in whole blood assay, and (2) has the cyclooxygenase-II  $IC_{50}$  value higher than  $0.2 \mu M$  in whole blood assay, shows excellent analgesic activity without causing undesired side effects, such as gastrointestinal disorders.

In the present invention, the more preferable analgesic agent is the one comprising the selective cyclooxygenase-I inhibitor, that (1) has a cyclooxygenase-II vs. cyclooxygenase-I  $IC_{50}$  values ratio higher than 50, more preferably 100, in whole blood assay, and that (2) has the cyclooxygenase-II  $IC_{50}$  value higher than  $0.5 \mu M$ , more preferably  $1.0 \mu M$ , in whole blood assay.

Accordingly, the analgesic agent of this invention are useful for treating or preventing acute or chronic pains caused by or associated with acute or chronic inflammations in human beings or animals by using administered systemically or topically.

Particularly, the following diseases are exemplified: pains caused by or associated with rheumatoid arthritis, osteoarthritis, lumbar rheumatism, rheumatoid spondylitis, gouty arthritis, juvenile arthritis, etc; lumbago; cervico-omo-brachial syndrome; scapulohumeral periartthritis; pain and tumescence after operation or injury; etc.

For therapeutic purpose, the analgesic agent of the present invention can be used in a form of pharmaceutical preparation suitable for oral, parenteral or external administration. The pharmaceutical preparations may be capsules, tablets, dragees, granules, inhalant, suppositories, solution, lotion, suspension, emulsion, ointment, gel, or the like.

The dosage of therapeutically effective amount of the analgesic agent will vary depending upon the age and condition

of each individual patient.

And further, the present application is concerning the followings.

5 An article of manufacture, comprising packaging material and the compound (I) identified in Claim 1 contained within said packaging material, wherein said the compound (I) is therapeutically effective for preventing or treating inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunity diseases, analgesic, thrombosis, cancer or  
10 neurodegenerative diseases, and wherein said packaging material comprises a label or a written material which indicates that said compound (I) can or should be used for preventing or treating inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunity diseases, analgesic,  
15 thrombosis, cancer or neurodegenerative diseases.

A commercial package comprising the pharmaceutical composition containing the compound (I) identified in Claim 1 and a written matter associated therewith, wherein the written  
20 matter states that the compound (I) can or should be used for preventing or treating inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunity diseases, analgesic, thrombosis, cancer or neurodegenerative diseases.

25

The patents, patent applications and publications cited herein are incorporated by reference.

30 The following Examples are given for the purpose of illustrating the present invention in detail.

#### Example 1

(1) A mixture of desoxyanisoin (8g, 31.2mol) and N,N-dimethylformamide dimethylacetal (9.3g, 78mmol) in dimethyl  
35 formamide (40ml) was stirred for 2 hours at 90°C.

The reaction mixture was evaporated under reduced pressure to afford crude 1-(N,N-dimethylamino)-2-(4-methoxyphenyl)-3-(4-methoxyphenyl)prop-1-en-3-one (10.72g) as a yellow solid. The crude solid was used for the next step without further purification.

(2) A mixture of 1-(N,N-dimethylamino)-2-(4-methoxyphenyl)-3-(4-methoxyphenyl)prop-1-en-3-one (10.64g, 31mmol) and 2-cyanoacetamide (2.92g, 34.7mmol) in N,N-dimethylformamide (80 ml) and methyl alcohol (3ml) was added to a slurry of NaH (2.73g, 68.2mmol : 60% in mineral oil) in N,N-dimethylformamide (40 ml) with cooling by an ice bath. (5 to 18 °C). The reaction mixture was stirred for 12hr at 80°C and cooled to room temperature. The resulting mixture was poured into 1 M KH<sub>2</sub>PO<sub>4</sub> (400ml), and filtered, washed with water (100ml) and dried in vacuo (60°C) to afford 1,2-dihydro-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-2-oxo-pyridine-3 carbonitrile (11.83g) as crystal.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, δ) : 3.80 (3H, s), 3.83 (3H, s), 6.81 (2H, d, J=8.8 Hz), 6.87 (2H, d, J=8.8 Hz), 6.98 (2H, d, J=8.8 Hz), 7.27 (2H, d, J=8.8 Hz), 7.92 (1H, s).

IR (KBr) : 2220, 1649, 1606, 1554, 1510, 1464, 1298, 1257, 1180, 1028 cm<sup>-1</sup>

Mass (ESI) : (M+H)<sup>+</sup> 333.1, (M+Na)<sup>+</sup> 355.2

(3) A mixture of 1,2-dihydro-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-2-oxo-pyridine-3-carbonitrile (2g, 4.85mmol), phosphorus oxychloride (2.26ml, 24.3mmol) and NEt<sub>3</sub> (0.676ml, 4.85mmol) was refluxed for 2hr. The resulting mixture was cooled to room temperature, concentrated under reduced pressure. The residue was dissolved in dichloromethane (10 ml) and 1N hydrochloric acid (10ml) (exothermic). The organic layer was separated and the aqueous layer was further extracted with dichloromethane (10 ml). The combined extracts were dried over MgSO<sub>4</sub> and concentrated under reduced pressure to afford deep brown solid (1.78g). The brown

solid was purified with column chromatography (Silica gel/  
Toluene) and triturated with ethyl acetate (3 ml) and concentrated  
to afford 2-chloro-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)  
pyridine-3-carbonitrile (1.39 g) as crystal.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.80 (3H, s), 3.83 (3H, s), 6.79 (2H, d,  $J=8.9$   
Hz), 6.87 (2H, d,  $J=8.8$  Hz), 7.10 (2H, d,  $J=8.8$  Hz), 7.37 (2H, d,  
 $J=8.9$  Hz), 7.90 (1H, s).

IR (KBr) : 2223, 1604, 1572, 1512, 1406, 1294, 1252, 1174, 1024  
 $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  351.20

### Example 2

2-Chloro-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-  
pyridine-3-carbonitrile (5.87g, 16.7mmol) was dissolved in  
dimethyl sulfoxide (64.6ml) at  $60^\circ\text{C}$  and then cooled to  $28^\circ\text{C}$  by water  
bath.  $\text{K}_2\text{CO}_3$  (6.94g, 50.2mmol) was added at small portion to the  
above solution under water bath cooling, successively 30%  $\text{H}_2\text{O}_2$  (5.87  
ml) was added to the reaction mixture (exothermic, 28 to  $36^\circ\text{C}$ ).  
The resulting mixture was stirred for 1hr under the same condition.  
The mixture was slowly poured into 1N hydrochloric acid (88.05  
ml, 15v) under ice bath (exothermic, 10 to  $25^\circ\text{C}$ ) to afford  
precipitates. The obtained precipitates were collected, washed  
with water (59 ml, 5v) fifth times and dried in vacuo to afford  
2-chloro-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-  
carboxamide (5.78 g).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.80 (3H, s), 3.82 (3H, s), 6.79 (2H, d,  $J=8.9$   
Hz), 6.84 (2H, d,  $J=8.8$  Hz), 7.13 (2H, d,  $J=8.8$  Hz), 7.38 (2H, d,  
 $J=8.9$  Hz), 8.26 (1H, s).

IR (KBr): 1673, 1603, 1579, 1512, 1392, 1292, 1252, 1176, 1024  
 $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  369.20

Example 3

To a mixture 2-chloro-5-(4-methoxyphenyl)-  
 6-(4-methoxyphenyl) -pyridine-3-carboxamide (4.02g, 10.9mmol)  
 and NEt<sub>3</sub> (15.2ml, 109mmol) in ethyl alcohol (20 ml) and THF (20  
 5 ml) was added Pd/C (800mg) . This mixture was hydrogenated for  
 2hr at 55°C and filtrated , washed with THF and ethyl alcohol  
 and concentrated to afford yellow solid . This crust was dissolved  
 in dichloromethane (40 ml, 10v) and water (40 ml, 10v) at 50°C.  
 The organic layer was separated, the aqueous layer was further  
 10 extracted with dichloromethane (20 ml), dried over MgSO<sub>4</sub> and  
 concentrated. The residue was triturated with ethyl acetate (16  
 ml, 4v) under reflux for 30 min, cooling to room temperature.  
 The resulting powder was collected, washed with ethyl acetate  
 (8 ml, 2v) twice and dried in vacuo to afford 5-(4-methoxyphenyl)  
 15 - 6-(4-methoxyphenyl) pyridine-3-carboxamide (2.94 g) as a  
 powder.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, δ) : 3.80(3H, s), 3.82(3H, s), 6.80(2H, d, J=8.8  
 Hz), 6.84(2H, d, J=8.8 Hz), 7.14(2H, d, J=8.8 Hz), 7.36(2H, d,  
 20 J=8.8 Hz), 8.12(1H, d, J=2.2 Hz), 8.99(1H, d, J=2.2 Hz).

IR (KBr) : 1682, 1510, 1383, 1292, 1252, 1176, 1028 cm<sup>-1</sup>

Mass (APCI) : (M+H)<sup>+</sup> 335.20

Example 4

25 A mixture of 5-(4-methoxyphenyl) - 6-(4-methoxyphenyl)  
 pyridine-3-carboxamide (2.89g, 8.64mmol) and phosphorus  
 oxychloride(14.2g, 92.9mmol) was stirred for 1hr under reflux .  
 The reaction mixture was cooled room temperature, concentrated,  
 codistilled with toluene and the residue was dissolved in ethyl  
 30 acetate (15 ml), washed with water (10 ml) three times, dried  
 over MgSO<sub>4</sub> and concentrated in vacuo to afford yellow solid. This  
 crust was purified by column chromatography (Silica gel/Toluene :  
 EtOAc = 10:1) . The obtained powder was recrystallized from n-butyl  
 alcohol (15 ml, 6v), collected by filtration, washed with n-butyl  
 35 alcohol (10 ml, 4v) twice and hexane (10 ml, 4v) twice and dried

in vacuo to afford 5- (4-methoxyphenyl)- 6-(4-methoxyphenyl)-pyridine-3-carbonitrile (2 g).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.80(3H, s), 3.83(3H, s), 6.80(2H, d,  $J=8.9$  Hz), 6.86(2H, d,  $J=8.8$  Hz), 7.11(2H, d,  $J=8.8$  Hz), 7.35(2H, d,  $J=8.9$  Hz), 7.90(1H, d,  $J=2.1$  Hz), 8.86(1H, d,  $J=2.1$  Hz).

IR (KBr) : 2223, 1581, 1508, 1423, 1292, 1248, 1173, 1022  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  317.40

mp: 107 - 108  $^{\circ}\text{C}$

#### Example 5

(1) A mixture of 1,2-dihydro-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-2-oxo-pyridine-3-carbonitrile (4g, 9.71mmol) and KOH (4.32 g) in ethylene glycol (16 ml, 4v), and water (6 ml, 1.5v) was heated at 160  $^{\circ}\text{C}$ . After stirring for over night, The reaction mixture was cooled to room temperature, and then poured into 1N hydrochloric acid (140 ml) to afford precipitates. The obtained precipitates were collected by filtration, washed with water twice and dried in vacuo to afford 1,2-dihydro-5-(4-methoxyphenyl)- 6-(4-methoxyphenyl)-2-oxo-pyridine-3-carboxylic acid (3.41g) as a crystal.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.81(3H, s), 3.86(3H, s), 6.82(2H, d,  $J=8.7$  Hz), 6.90(2H, d,  $J=8.8$  Hz), 7.05(2H, d,  $J=8.7$  Hz), 7.29(2H, d,  $J=8.8$  Hz), 8.62(1H, s), 12.71(1H, brs), 13.61(1H, brs).

IR (KBr) : 1728, 1616, 1552, 1506, 1450, 1257, 1176  $\text{cm}^{-1}$

Mass (ESI) :  $(\text{M}+\text{Na})^+$  374.2

(2) 1,2-Dihydro-5-(4-methoxyphenyl)- 6-(4-methoxyphenyl)-2-oxo-pyridine-3-carboxylic acid (0.8g, 2.28mmol) was heated at 210 $^{\circ}\text{C}$  in quinoline (5ml). After stirring for over night, ethyl acetate and 1N-hydrochloric acid were added to the reaction mixture. The precipitates were collected by filtration, washed with 1N-hydrochloric acid and ethyl acetate and dried in vacuo to afford 1,2-dihydro-5-(4-methoxyphenyl)- 6-(4-methoxyphenyl)

-2-oxo-pyridine (1.36 g). The crude solid was used for the next step without further purification.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.79(3H, s), 3.81(3H, s), 6.62(1H, d,  $J=9.3$  Hz), 6.78(2H, d,  $J=8.6$  Hz), 6.83(2H, d,  $J=8.6$  Hz), 6.99(2H, d,  $J=8.6$  Hz), 7.18(2H, d,  $J=8.6$  Hz), 7.56(1H, d,  $J=9.3$  Hz).

Mass (APCI) :  $(\text{M}+\text{H})^+$  308.27

(3) To a solution of 1,2-dihydro-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-2-oxo-pyridine (0.9g, 1.6mmol) in pyridine was added trifluoromethanesulfonic acid anhydride (0.808ml, 4.8mmol) and warmed for 1hr at  $60^\circ\text{C}$ . The reaction mixture was concentrated and then purified by column chromatography (Silica gel / 40%dichloromethane/hexane) to afford trifluoromethanesulfonic acid 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-2-yl ester (374mg)

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.80(3H, s), 3.82(3H, s), 6.78(2H, d,  $J=8.9$  Hz), 6.86(2H, d,  $J=8.8$  Hz), 7.00 - 7.20(3H, m), 7.33(2H, d,  $J=8.9$  Hz), 7.80(1H, d,  $J=8.2$  Hz).

IR (KBr) : 1603, 1585, 1514, 1446, 1417, 1250, 1174, 1128  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  439.87

(4) A mixture of trifluoromethanesulfonic acid 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-2-yl ester (80mg, 0.182mmol), KCN (35.6mg, 0.546mmol), LiCl (23.2mg, 0.546mmol), 18crown6 (14mg, 0.3eq) and palladium tetrakis(triphenylphosphine) (42.1mg, 0.0364mmol) in toluene (5ml) was heated for 15hr at  $100^\circ\text{C}$ . The reaction mixture was cooled to room temperature, and then extracted with EtOAc at several times. The organic layer was washed with water, dried over  $\text{MgSO}_4$  and concentrated. The residue was purified by thin layer chromatography (20%Hexane in EtOAc) to afford 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-2-carbonitrile (27mg) as crystal.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.80 (3H, s), 3.82 (3H, s), 6.81 (2H, d,  $J=8.8$  Hz), 6.85 (2H, d,  $J=8.8$  Hz), 7.12 (2H, d,  $J=8.8$  Hz), 7.33 (2H, d,  $J=8.8$  Hz), 7.63 (1H, d,  $J=7.9$  Hz), 7.76 (1H, d,  $J=7.9$  Hz).

IR (KBr) : 2233, 1512, 1246, 1174, 1028  $\text{cm}^{-1}$

5 Mass (APCI) :  $(\text{M}+\text{H})^+$  317.33

#### Example 6

A mixture of 2-chloro-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-carbonitrile (124mg, 0.353mmol) and 28%NaOMe in methyl alcohol (5ml) in N,N-dimethylformamide was refluxed for 10 2hr. The reaction mixture was cooled and concentrated under reduced pressure. The residue was diluted with ethyl acetate, washed with water twice and brine, dried over  $\text{MgSO}_4$  and concentrated (0.14g). The crude product was purified by thin layer chromatography 15 (20%Hexane in EtOAc) to afford 2-methoxy-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-carbonitrile (0.06 g) as a powder.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.80 (3H, s), 3.82 (3H, s), 4.13 (3H, s), 6.78 (2H, d,  $J=8.8$  Hz), 6.82 (2H, d,  $J=8.8$  Hz), 7.07 (2H, d,  $J=8.8$  Hz), 7.39 (2H, d,  $J=8.8$  Hz), 7.81 (1H, s). 20

IR (KBr) : 2225, 1591, 1462, 1398, 1250, 1173, 1028  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  347.40

#### Example 7

(1) To a solution of 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-carbonitrile (0.13g, 0.41mmol) in toluene (5ml) under nitrogen atmosphere was added DIBAL (0.82 ml : 1M in toluene) at -78°C and stirred for 2hr at room temperature. The reaction mixture was quenched by 1N hydrochloric acid, basified by sat.  $\text{NaHCO}_3$  30 aq., extracted with ethyl acetate twice, dried over  $\text{MgSO}_4$  and concentrated.

The residue was purified by column chromatography (Silica gel / 40%EtOAc / hexane) to afford 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-carboaldehyde (62mg).



$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.81(3H, s), 3.83(3H, s), 6.81(2H, d,  $J=8.8$  Hz), 6.86(2H, d,  $J=8.8$  Hz), 7.15(2H, d,  $J=8.8$  Hz), 7.40(2H, d,  $J=8.8$  Hz), 8.12(1H, d,  $J=2.1$  Hz), 9.05(1H, d,  $J=2.1$  Hz), 10.16(1H, s).

5 IR (KBr) : 1695, 1583, 1512, 1248, 1174, 1028  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  320.33

(2) To a solution of 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-carboaldehyde(57mg, 0.178mmol) in dichloromethane  
10 (5ml) was added (diethylamino)sulfur trifluoride (86.3mg, 0.535mmol) at  $0^\circ\text{C}$ . The reaction mixture was stirred for over night at room temperature. The resulting mixture concentrated and purified by thin layer chromatography (30% Hexane in EtOAc) to afford 3-difluoromethyl-5-(4-methoxyphenyl)-6-  
15 (4-methoxyphenyl)-pyridine(26 mg).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.80(3H, s), 3.82(3H, s), 6.40 -7.10(5H, m), 7.13(2H, d,  $J=8.8$  Hz), 7.34(2H, d,  $J=8.8$  Hz), 7.80(1H, s), 8.73(1H, s).

20 IR (KBr):1604, 1512, 1452, 1427, 1365, 1252, 1176, 1088, 1032  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  342.33

#### Example 8

25 (1) To a solution of 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-carbonitrile(400mg, 1.26mmol) in ethyl alcohol(10ml) and c-hydrochloric acid(600ul) was added 10%Pd/C (50%wet, 80mg). The reaction mixture was hydrogenated for 2hr at  $55^\circ\text{C}$ . The resulting mixture was filtered and concentrated. The  
30 residue was resolved in ethyl acetate and 1N hydrochloric acid aq.. The aqueous layer was separated and the organic layer was further extracted with 1N hydrochloric acid aq.. The combined hydrochloric acid layer was basified by 1N NaOH aq. and extracted with dichloromethane three times. The combined organic layer was  
35 dried over  $\text{MgSO}_4$  and concentrated. The residue was purified by

column chromatography (Silica gel /15% methyl alcohol/CH<sub>3</sub>Cl) to afford 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-methyl-amine (305mg).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, δ) : 3.79 (3H, s), 3.81 (3H, s), 3.97 (2H, s), 6.78 (2H, d, J=8.9 Hz), 6.82 (2H, d, J=8.9 Hz), 7.12 (2H, d, J=8.9 Hz), 7.30 (2H, d, J=8.9 Hz), 7.65 (1H, d, J=2.2 Hz), 8.57 (1H, d, J=2.2 Hz).  
IR (neat) : 1608, 1512, 1292, 1242, 1178, 1032 cm<sup>-1</sup>  
Mass (APCI) : (M+H)<sup>+</sup> 321.33

(2) To a mixture of 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-methylamine (139mg, 0.434mmol) and 35% HCHO aq. (12.6M, 344ul) in dichloromethane (5 ml) and methyl alcohol (2 ml) was added NaBH(OAc)<sub>3</sub> (552 mg, 2.6 mmol) at room temperature and then stirred for 30min. The reaction was quenched with water, extracted with dichloromethane twice, dried over MgSO<sub>4</sub> and concentrated. The residue was purified by thin layer chromatography (10% methyl alcohol in dichloromethane) to afford an oil.

This oil was dissolved in dichloromethane (5 ml) was treated with 4N hydrochloric acid in ethyl acetate (1 ml) (suspension) and concentrated. The hydrochloric acid salts were triturated with dichloromethane and iso-propyl ether and concentrated to afford 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-N,N-dimethylmethylamine hydrochloride (98 mg).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, δ) : 3.00 (6H, s), 3.81 (3H, s), 3.85 (3H, s), 4.60 - 5.00 (2H, m), 6.70 - 7.80 (8H, m), 9.37 (1H, s), 9.52 (1H, s).  
IR (KBr) : 1606, 1510, 1252, 1182, 1024 cm<sup>-1</sup>  
Mass (APCI) : (M+H)<sup>+</sup> (free) 349.27

### Example 9

A mixture of 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-methylamine (96mg, 0.3mmol) and methylisocyanate (25.3mg, 0.449mmol) in THF (5 ml) and methyl alcohol (1 ml) was stirred for 1hr at room temperature. The reaction was concentrated

and the residue was purified by thin layer chromatography (10% methyl alcohol in dichloromethane) to afford an oil. This oil was treated with 4N hydrochloric acid in ethyl acetate and concentrated. The residue was triturated with  
 5 dichloromethane/iso-propyl ether to afford a pale yellow powder N-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-3-methylpyridine-N'-methylurea hydrochloride (95 mg).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, δ) : 2.75 (3H, s), 3.82 (6H, s), 4.64 (2H, s), 6.85 (2H, d, J=3.1 Hz), 6.89 (2H, d, J=3.1 Hz), 7.10 (2H, d, J=8.6 Hz), 7.38 (2H, d, J=8.6 Hz), 8.35 (1H, brs), 8.95 (1H, brs).

IR (KBr) : 2058, 1652, 1606, 1510, 1460, 1302, 1255, 1180 cm<sup>-1</sup>

Mass (APCI) : (M+H)<sup>+</sup> (free) 378.00

#### 15 Example 10

To a solution of 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-carbonitrile (24mg, 0.076mmol) in ethyl alcohol (5ml) and c-hydrochloric acid (40 μl) was added 10% Pd/C (50% wet, 10mg). The reaction mixture was hydrogenated for 2.5hr at 55°C. The  
 20 resulting mixture was filtrated and concentrated to afford 5-(4-methoxyphenyl)-6-(4-methoxyphenyl)-pyridine-3-methylamine hydrochloride (31 mg) as pale yellow powder.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, δ) : 3.68 (3H, s), 3.71 (3H, s), 3.84 (2H, s), 6.40  
 25 - 8.00 (10H, m).

IR (KBr) : 1605, 1510, 1257, 1180, 1022 cm<sup>-1</sup>

Mass (APCI) : (M+H)<sup>+</sup> 321.27

#### 30 Example 11

(1) A mixture of 3-bromo-5-methyl-2-aminopyridine (4.0g, 21.4mmol), 4-methoxybenzeneboronic acid (3.9g, 25.7mmol) and palladium tetrakis(triphenylphosphine) (247mg, 0.214mmol) in benzene (20ml) -ethyl alcohol (20ml) -2M Na<sub>2</sub>CO<sub>3</sub> (24ml) was refluxed  
 35 for 16hr. The reaction mixture was diluted with ethyl acetate

and water and the organic layer was separated. The aqueous layer was further extracted with ethyl acetate. The combined organic layer was dried over  $\text{MgSO}_4$  and concentrated. The residue was purified by column chromatography ( Silica gel / 20-70% ethyl acetate/hexane ) to afford 5-methyl-3-(4-methoxyphenyl)-2-aminopyridine (4.71 g)

$^1\text{H NMR}$  ( $\text{CDCl}_3$ ,  $\delta$ ) : 2.22 (3H, s), 3.85 (3H, s), 4.44 (2H, brs), 6.98 (2H, d,  $J=8.8$  Hz), 7.18 (1H, d,  $J=2.1$  Hz), 7.37 (2H, d,  $J=8.8$  Hz), 7.88 (1H, s,  $J=2.1$  Hz).

IR (KBr) : 1627, 1610, 1568, 1508, 1464, 1406, 1284, 1244, 1173, 1026  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  215.27

(2) To a mixture of 5-methyl-3-(4-methoxyphenyl)-2-aminopyridine (1g, 4.57mmol) in ethyl alcohol (10ml) - 1.88M  $\text{H}_2\text{SO}_4$  solution was added n-butyl alcohol (10 ml),  $\text{NaNO}_2$  (5.15g, 74.7mmol) and then the resulting mixture was stirred for 4h at  $65^\circ\text{C}$ . The mixture was diluted with ethyl acetate and water and the organic layer was separated. The aqueous layer was further extracted with ethyl acetate. The combined organic layer was dried over  $\text{MgSO}_4$  and concentrated. The residue was triturated with ethyl acetate and iso-propyl ether, collected by filtration, washed with iso-propyl ether and dried in vacuo to afford 5-methyl-3-(4-methoxyphenyl)-2-hydroxypyridine (0.55g) as pale orange powder.

$^1\text{H NMR}$  ( $\text{CDCl}_3$ ,  $\delta$ ) : 2.13 (3H, s), 3.85 (3H, s), 6.96 (2H, d,  $J=8.8$  Hz), 7.12 (1H, d,  $J=2.4$  Hz), 7.41 (1H, d,  $J=2.4$  Hz), 7.68 (2H, d,  $J=8.8$  Hz), 12.64 (1H, brs).

IR (KBr) : 1657, 1562, 1510, 1290, 1246, 1173, 1024  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  216.20

(3) Trifluoromethanesulfonic acid 3-methyl 3-(4-methoxyphenyl)pyridine-2-yl ester was prepared from 5-methyl-

3-(4-methoxyphenyl)-2-hydroxypyridine by the similar method as that described for Example 5-(3).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 2.41(3H, s), 3.87(3H, s), 7.00(2H, d,  $J=8.8$  Hz), 7.40(2H, d,  $J=8.8$  Hz), 7.66(1H, d,  $J=2.0$  Hz), 8.11(1H, d,  $J=2.0$  Hz).

IR (neat) : 1610, 1516, 1414, 1252, 1215, 1140, 1038  $\text{cm}^{-1}$

Mass (ESI) :  $(\text{M}+\text{H})^+$  348.1,  $(\text{M}+\text{Na})^+$  370.1

(4) 2-Methyl-5-(4-methoxyphenyl)-6-(4-methoxyphenyl)pyridine hydrochloride was prepared from trifluoromethanesulfonic acid 3-methyl 3-(4-methoxyphenyl) pyridine-2-yl ester by the similar method as that described for Example 11-(1).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 2.56(3H, s), 3.83(6H, s), 6.87(2H, d,  $J=2.9$  Hz), 6.91(2H, d,  $J=2.9$  Hz), 7.11(2H, d,  $J=8.8$  Hz), 7.46(2H, d,  $J=8.8$  Hz), 8.06(1H, s), 8.73(1H, s).

IR (KBr) : 2089, 1606, 1508, 1460, 1255, 1178, 1024  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  (free) 306.20

## Example 12

(1) 2-Benzyloxy-5-chloro-3-(4-methoxyphenyl) pyridine was prepared from 2-benzyloxy-3-bromo-5-chloropyridine by the similar method as that described for Example 11-(1)

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.85(3H, s), 5.44(2H, s), 6.95(2H, d,  $J=8.9$  Hz), 7.20 - 7.50(5H, m), 7.53(2H, d,  $J=8.9$  Hz), 7.60(1H, d,  $J=2.6$  Hz), 8.05(1H, d,  $J=2.6$  Hz).

IR (KBr) : 1608, 1510, 1435, 1362, 1302, 1244, 1174, 1032  $\text{cm}^{-1}$

Mass (APCI) :  $(\text{M}+\text{H})^+$  326.13

(2) A mixture of 2-benzyloxy-5-chloro-3-(4-methoxyphenyl) pyridine in 6N hydrochloric acid(10.5ml) and ethyl alcohol/toluene(1/1, 10.5ml) was refluxed for 2h. The reaction mixture was cooled to room temperature and diluted with ethyl

acetate and water. The organic layer was separated and the aqueous layer was further extracted with dichloromethane. The crystal was collected by filtration (5-chloro-3-(4-methoxyphenyl)-2-hydroxypyridine:0.32g), washed with ethyl acetate and dried in vacuo. The combined filtrate was dried over  $\text{MgSO}_4$  and concentrated. The residual solid was triturated with iso-propyl ether, collected by filtration, washed with iso-propyl ether and dried in vacuo to afford 5-chloro-3-(4-methoxyphenyl)-2-hydroxypyridine (0.98 g).

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ,  $\delta$ ) : 3.78 (3H, s), 6.96 (2H, d,  $J=8.9$  Hz), 7.56 (1H, d,  $J=2.9$  Hz), 7.62 (1H, d,  $J=2.9$  Hz), 7.72 (2H, d,  $J=8.9$  Hz), 11.98 (1H, brs).

IR (KBr) : 1651, 1604, 1510, 1468, 1250, 1178, 1022  $\text{cm}^{-1}$

(3) Trifluoromethanesulfonic acid 5-chloro-3-(4-methoxyphenyl)pyridine-2-yl ester was prepared from 5-chloro-3-(4-methoxyphenyl)-2-hydroxypyridine by the similar method as that described for Example 5-(3).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.87 (3H, s), 7.02 (2H, d,  $J=8.8$  Hz), 7.41 (2H, d,  $J=8.8$  Hz), 7.85 (1H, d,  $J=2.5$  Hz), 8.25 (1H, d,  $J=2.5$  Hz).

IR (neat) : 1610, 1516, 1421, 1252, 1217, 1140, 1034  $\text{cm}^{-1}$

Mass (ESI) :  $(\text{M}+\text{H})^+$  368.0,  $(\text{M}+\text{Na})^+$  390.1

(4) 5-Chloro-2-(4-methoxyphenyl)-3-(4-methoxyphenyl)pyridine was prepared from trifluoromethanesulfonic acid 5-chloro-3-(4-methoxyphenyl)pyridine-2-yl ester (XVII) by the similar method as that described for Example 11-(1).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ,  $\delta$ ) : 3.79 (3H, s), 3.81 (3H, s), 6.78 (2H, d,  $J=8.8$  Hz), 6.83 (2H, d,  $J=8.8$  Hz), 7.10 (2H, d,  $J=8.8$  Hz), 7.28 (2H, d,  $J=8.8$  Hz), 7.66 (1H, d,  $J=2.4$  Hz), 8.57 (1H, d,  $J=2.4$  Hz).

IR (neat) : 1608, 1512, 1460, 1429, 1246, 1178, 1115, 1030  $\text{cm}^{-1}$

Mass (ESI) : (M+H)<sup>+</sup> 326.3, (M+Na)<sup>+</sup> 348.1

Example 13

(1) To a suspension of 6-amino-nicotinamide (4.45 g, 32.4 mmol) in acetic acid (100 ml) was added dropwise bromine (1.84 ml, 35.7 mmol) at room temperature. The mixture was stirred at 55°C for 3 hours. The resulting solution was added to 750 ml of 3N aqueous sodium hydroxide. The resulting sediment was filtered, washed with water and was dried over to afford 6-amino-5-bromonicotinamide. This was used for the next step without further purification.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, δ); 6.77 (2H, br), 7.19 (1H, br), 7.77 (1H, br), 8.15 (1H, d, J=2.0 Hz), 8.47 (1H, d, J=2.0 Hz),

MASS (APCI) ; 216 (M+H)<sup>+</sup>

(2) To the solution of 6-amino-5-bromonicotinamide (6.67 g, 30.9 mmol), 4-methoxybenzeneboronic acid (5.63 g, 37 mmol) and tetrakis(triphenylphosphin) -palladium(0) (1.78 g, 1.54 mmol) in ethylene glycol dimethylether (60 ml) was added 2M aqueous sodium carbonate (92 ml). And this was stirred for 15 hours at 100°C. After the mixture was cooled to room temperature, this was diluted with 1N aqueous sodium hydroxide. The reaction mixture was extracted with ethyl acetate. The extract was washed with brine, dried over magnesium sulfate and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate (1:1) as an eluent to afford 6-amino-5-(4-methoxyphenyl)-nicotinamide (5.97 g, white solid).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, δ); 3.80 (3H, s), 6.09 (2H, s), 7.04 (3H, m, J=8.8 Hz), 7.38 (3H, d, J=8.8 Hz), 7.75 (2H, m), 8.47 (1H, d, J=1.9 Hz),

MASS (ESI) ; 266 (M + Na)<sup>+</sup>

(3) 6-Amino-5-(4-methoxyphenyl)nicotinamide (700 mg, 2.88 mg)

was dissolved into acetone (15 ml) and 1.88 M H<sub>2</sub>SO<sub>4</sub> (15 ml) . To this solution, 5 M NaNO<sub>2</sub> (5.8 ml) was added dropwise under ice cooling. And the resultant solution was stirred for 5h. (Bubbled, Brown gas). Another NaNO<sub>2</sub> (567 mg, in 2 ml water) was added at 0°C and was stirred for 3h. Sediment was filtrated, washed with water and dried up to give 6-hydroxy-5-(4-methoxyphenyl)-nicotinamide (543 mg). This was used for the next step without further purification.

(4) To the solution of 6-hydroxy-5-(4-methoxyphenyl)-nicotinamide (1.8 g, 7.37 mmol) and triethylamine hydrochloride (7.1 g, 51.6 mmol) in toluene was added phosphorous chloride. After being stirred for 12 hours at 110°C, the reaction mixture was poured into water, extracted with ethyl acetate. The extract was washed with brine, dried over magnesium sulfate and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography with a mixture of hexane and ethyl acetate (2:1) as an eluent to afford 6-chloro-5-(4-methoxyphenyl)nicotinonitrile.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, δ); 3.88 (3H, s), 7.01 (2H, d, J=9.7 Hz), 7.39 (2H, d, J=9.7 Hz), 7.90 (1H, d, J=2.3 Hz), 8.62 (1H, d, J=2.3 Hz),

IR (cm<sup>-1</sup>); 1693, 1617, 1515, 1380, 1251, 1186, 1095, 1025, 827

MASS (APCI) : 245 (M+H)+

(5) To the solution of 6-chloro-5-(4-methoxyphenyl)-nicotinonitrile (220 mg, 0.899 mmol) in dimethoxyethane (5 ml), 6-methoxy-3-pyridinylboronic acid (344 mg, 2.25 mmol), tetrakis(triphenylphosphin)palladium (31.2 mg, 0.027 mmol) and 2M Na<sub>2</sub>CO<sub>3</sub> (1.8 ml, 36mmol) was added. The reaction mixture was stirred at 80°C for 12 h.

This was diluted with ethyl acetate, washed with 0.1N hydrochloric acid and brine. After subjecting an extraction with ethyl acetate,



a purification by silica gel column chromatography was carried out with hexane/ ethyl acetate (5-3/1) as an eluent. And 5-(4-methoxyphenyl)-6-(6-methoxypyridine-3-yl)-nicotinonitrile (colorless crystal needle) was obtained by recrystallization from ethyl alcohol.

IR (KBr,  $\text{cm}^{-1}$ ) ; 2225, 1602, 1575, 1504, 1438, 1400, 1371, 1309, 1292, 1245, 1174, 1118, 1064, 1014, 939, 827, 786

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ,  $\delta$ ) ; 3.83 (3H, s), 3.93 (3H, s), 6.64 (1H, d,  $J=8.7$  Hz), 6.88 (2H, d,  $J=9$  Hz), 7.1 (2H, d,  $J=9$  Hz), 7.6 (1H, dd,  $J=8.725$  Hz), 7.92 (1H, d,  $J=2.1$  Hz), 8.22 (1H, d,  $J=2.5$  Hz), 8.88 (1H, d,  $J=2.1$  Hz),

MASS (APCI) ; 318 ( $\text{M}+\text{H}$ )+

#### Example 14

A mixture of the compound obtained in a similar manner to that of Example 1-(3) (7.02g, 20mmol) and zinc powder (5.26g) in a mixture of acetic acid (80mL) and  $\text{N,N}$ -dimethylformamide (40mL) was stirred at  $55^\circ\text{C}$  for 8 h.

Then the reaction mixture was stirred at room temperature for 14 h. Zinc salt was removed by filtration. To the filtrate was added toluene (200mL) and water (100mL). The layers were separated and the aqueous layer was re-extracted with toluene (100 mL). The combined organic layer was washed with water (100 mL) and 10% brine (100 mL) respectively.

The organic layer was evaporated and the residual waxy oil was treated with 50% ethyl alcohol in water. The precipitate was collected and dried under reduced pressure to afford the same compound as Example 4 (5.37g, 85% yield) as a yellowish solid.

















$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ,  $\delta$ ) ; 3.80 (s, 3H), 3.82 (s, 3H), 4.60 (s, 2H), 6.79 (dd, 2H,  $J=8.9, 2.0$ ), 6.86 (dd, 2H,  $J=8.7, 2.0$ ), 7.11 (dd, 2H,  $J=8.8, 2.1$ ), 7.35 (dd, 2H,  $J=8.8, 2.1$ ), 7.89 (d, 1H,  $J=2.1$ ), 8.85 (d,

$1\text{H}, J=2.0);$ 

 MS (EI);  $m/z$  317.3 (M+H)<sup>+</sup>


5 The chemical structures of the compounds produced by the above Examples are listed in the following Table 2.

Table 2

Example No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
1-(3)	-CN	-Cl		-OCH <sub>3</sub>
2	-CONH <sub>2</sub>	-Cl		-OCH <sub>3</sub>
3	-CONH <sub>2</sub>	-H		-OCH <sub>3</sub>
4	-CN	-H		-OCH <sub>3</sub>
5-(4)	-H	-CN		-OCH <sub>3</sub>
6	-CN	-OCH <sub>3</sub>		-OCH <sub>3</sub>
7-(1)	-CHO	-H		-OCH <sub>3</sub>
7-(2)	-CHF <sub>2</sub>	-H		-OCH <sub>3</sub>
8-(1)	-CH <sub>2</sub> NH <sub>2</sub>	-H		-OCH <sub>3</sub>
8-(2)	-CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	-H		-OCH <sub>3</sub>
9	-CH <sub>2</sub> NH-CO-NHCH <sub>3</sub>	-H		-OCH <sub>3</sub>
10	-CH <sub>2</sub> NH <sub>2</sub>	-H		-OCH <sub>3</sub>
11-(4)	-CH <sub>3</sub>	-H		-OCH <sub>3</sub>
12-(4)	-Cl	-H		-OCH <sub>3</sub>
13-(5)	-CN	-H		-OCH <sub>3</sub>
14	-CN	-H		-OCH <sub>3</sub>

## C L A I M S

1. A compound of the formula (I):



(I)

wherein R<sup>1</sup> is hydrogen, halogen, carbamoyl, cyano, formyl, or lower alkyl optionally substituted with halogen, amino or a protected amino; R<sup>2</sup> is hydrogen, halogen, cyano or lower alkoxy; R<sup>3</sup> is phenyl or pyridyl, each of which is substituted with lower alkoxy; and R<sup>4</sup> is lower alkoxy; provided that either R<sup>1</sup> or R<sup>2</sup> is hydrogen, then the other is other than hydrogen, or its salts.

2. The compound of Claim 1, in which  
R<sup>1</sup> is halogen, carbamoyl, cyano, formyl, or lower alkyl optionally substituted with halogen, amino or a protected amino;  
R<sup>2</sup> is hydrogen;  
R<sup>3</sup> is phenyl substituted with lower alkoxy, or pyridyl substituted with lower alkoxy; and  
R<sup>4</sup> is lower alkoxy.
3. The compound of Claim 1, in which  
R<sup>1</sup> is hydrogen;  
R<sup>2</sup> is halogen, cyano or lower alkoxy;  
R<sup>3</sup> is phenyl substituted with lower alkoxy; and  
R<sup>4</sup> is lower alkoxy.

4. The compound of Claim 1, in which

$R^1$  is cyano, or lower alkyl optionally substituted with  
halogen, amino or a protected amino;

$R^2$  is halogen, cyano or lower alkoxy;

5  $R^3$  is phenyl substituted with lower alkoxy; and

$R^4$  is lower alkoxy.

10 5. A pharmaceutical composition comprising the compound (I)  
or its salts of Claim 1, as an active ingredient, in association  
with a pharmaceutically non-toxic carrier or excipient.

15 6. The composition of Claim 5, which is for treating and/or  
preventing inflammatory conditions, various pains, collagen  
diseases, autoimmune diseases, various immunity diseases,  
analgesic, thrombosis, cancer or neurodegenerative diseases

7. A compound of Claim 1 for use as a medicament.

20 8. A method for treatment and/or prevention of inflammatory  
conditions, various pains, collagen diseases, autoimmune diseases,  
various immunity diseases, analgesic, thrombosis, cancer or  
neurodegenerative diseases which comprises administering an  
effective amount of the compound (I) or its salts of Claim 1 to  
25 human beings or animals.

30 9. A use of the compound of Claim 1 for the manufacture of a  
medicament for treatment and/or prevention of inflammatory  
conditions, various pains, collagen diseases, autoimmune diseases,  
various immunity diseases, analgesic, thrombosis, cancer or  
neurodegenerative diseases in human beings or animals.

35 10. An analgesic agent comprising a selective cyclooxygenase-I  
inhibitor, a cyclooxygenase-II vs. cyclooxygenase-I  $IC_{50}$  values  
ratio of which is higher than 30 in a whole blood assay and

cyclooxygenase-II  $IC_{50}$  value thereof is higher than  $0.2 \mu M$  in a whole blood assay.

11. The analgesic agent of Claim 10, which is usable for treating and/or preventing pains caused by or associated with acute or chronic inflammations without causing gastrointestinal disorders.

12. The analgesic agent of Claim 10, which is usable for treating or preventing pains caused by or associated with rheumatoid arthritis, osteoarthritis, lumbar rheumatism, rheumatoid spondylitis, gouty arthritis, or juvenile arthritis; lumbago; cervico-omo-brachial syndrome; scapulohumeral periarthrititis; pain and tumescence after operation or injury without causing gastrointestinal disorders.

13. The analgesic agent of Claim 10, in which the  $IC_{50}$  values ratio is higher than 50.

14. The analgesic agent of Claim 10, in which the  $IC_{50}$  values ratio is higher than 100.

15. The analgesic agent of Claim 10, in which the cyclooxygenase-II  $IC_{50}$  value of the selective cyclooxygenase-I inhibitor is higher than  $0.5 \mu M$ .

16. A method for treating and/or preventing inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunity diseases, analgesic, thrombosis, cancer or neurodegenerative diseases, which comprises administering an effective amount of a selective cyclooxygenase-I inhibitor, which is characterized by a cyclooxygenase-II vs. cyclooxygenase-I  $IC_{50}$  values ratio of which is higher than 30 in a whole blood assay, and cyclooxygenase-II  $IC_{50}$  value thereof is higher than  $0.2 \mu M$  in a whole blood assay,

to human beings or animals.

17. A use of a selective cyclooxygenase-I inhibitor,  
which is characterized by a cyclooxygenase-II vs.  
5 cyclooxygenase-I IC<sub>50</sub> values ratio of which is higher  
than 30 in a whole blood assay and cyclooxygenase-II  
IC<sub>50</sub> value thereof is higher than 0.2  $\mu$ M in a whole blood  
assay,

10 for the manufacture of a medicament for treatment and/or  
prevention of inflammatory conditions, various pains, collagen  
diseases, autoimmune diseases, various immunity diseases,  
analgesic, thrombosis, cancer or neurodegenerative diseases.

18. A method for selecting a selective cyclooxygenase-I  
15 inhibitor, which lacks gastrointestinal disorders, by assessing  
whether cyclooxygenase-II vs. cyclooxygenase-I IC<sub>50</sub> values ratio  
is higher than 30 in a whole blood assay and whether the  
cyclooxygenase-II IC<sub>50</sub> value thereof is higher than 0.2  $\mu$ M in whole  
blood assay.

20

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 01/11241

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D213/85 C07D213/82 C07D213/84 C07D213/61 C07D213/48  
 C07D213/26 C07D213/38 C07D213/40 A61K31/4418 A61P25/00  
 A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 24584 A (SEARLE & CO ;WEIER RICHARD M (US); LEE LEN F (US); PARTIS RICHARD) 15 August 1996 (1996-08-15) cited in the application claims 8,10-32  -/-	1,5, 7-10, 16-18

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the International search

17 May 2002

Date of mailing of the International search report

04/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Bosma, P

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 01/11241

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; KONNO, SHOETSU ET AL: "Studies on as-triazine derivatives. XIX. Synthesis of 2,3-diarylpyrazine and 2,3-diarylpyridine derivatives as blood platelet aggregation inhibitors" retrieved from STN Database accession no. 118:234009 XP002199240 abstract & YAKUGAKU ZASSHI (1993), 113(1), 40-52 , -----	1,5,7
X	MARCOUX, JEAN-FRANCOIS ET AL: "Annulation of Ketones with Vinamidinium Hexafluorophosphate Salts: An Efficient Preparation of Trisubstituted Pyridines" ORGANIC LETTERS, vol. 2 , no. 15, 2000, pages 2339-2341, XP002199239 the whole document -----	1
A	WO 98 03484 A (GAUTHIER JACQUES YVES ;MERCK FROSST CANADA INC (CA); DUBE DANIEL ( ) 29 January 1998 (1998-01-29) cited in the application the whole document -----	1,5, 7-10, 16-18
A	WO 92 02513 A (FUJISAWA PHARMACEUTICAL CO., LTD., JAPAN) 20 February 1992 (1992-02-20) the whole document -----	1-18



## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 01/11241

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9624584	A	15-08-1996	US 5686470 A	11-11-1997
			AT 198327 T	15-01-2001
			AU 4859396 A	27-08-1996
			DE 69611354 D1	01-02-2001
			DE 69611354 T2	07-06-2001
			DK 808304 T3	29-01-2001
			EP 0808304 A1	26-11-1997
			ES 2154398 T3	01-04-2001
			PT 808304 T	29-06-2001
			WO 9624584 A1	15-08-1996
			US 5916905 A	29-06-1999
WO 9803484	A	29-01-1998	AU 723179 B2	17-08-2000
			AU 3331997 A	10-02-1998
			BG 103179 A	30-11-1999
			CA 2260016 A1	29-01-1998
			WO 9803484 A1	29-01-1998
			CZ 9900130 A3	16-06-1999
			EE 9900018 A	16-08-1999
			EP 0912518 A1	06-05-1999
			HR 970389 A1	30-06-1998
			JP 11514008 T	30-11-1999
			JP 3251945 B2	28-01-2002
			JP 2002080453 A	19-03-2002
			NO 990191 A	16-03-1999
			NZ 333230 A	25-08-2000
			PL 330995 A1	21-06-1999
			SK 3699 A3	10-12-1999
			TR 9900046 T2	21-04-1999
			BR 9710372 A	17-08-1999
			HU 9903974 A2	28-03-2000
			US 6001843 A	14-12-1999
			US 6071936 A	06-06-2000
			US 5861419 A	19-01-1999
WO 9202513	A	20-02-1992	WO 9202513 A1	20-02-1992
			JP 6501926 T	03-03-1994

**This Page Blank (uspto)**